

CERTAIN QUADRUPLE SERIES EQUATIONS INVOLVING LAGUERRE POLYNOMIALS

By

Omkar Lal Shrivastava¹ and Kuldeep Narain²

¹Department of Mathematics

Government Kamladevi Rathi Girls Postgraduate College, Rajnandgaon-491441, Chhattisgarh, India

Email:omkarlal@gmail.com

²Department of Mathematics

Kymore Science College, Kymore-483880, Madhya Pradesh, India

Email:kuldeepnarain2009@gmail.com

(Received : November 30, 2020 ; Revised: December 24, 2020)

DOI: <https://doi.org/10.58250/jnanabha.2020.50232>

Abstract

Srivastava ([13], [15]) has solved dual series equations involving Bateman- k functions and Jacobi polynomials. Srivastava [16] has obtained more results for the Konhauser-biorhogonal set. Lowndes ([3], [4]), Srivastava [12], Lowndes and Srivastava [5], Srivastava[14], Srivastava and Panda [17] have obtained the solution of dual series equations involving Jacobi and Laguerre polynomials and also solved triple series equations involving Laguerre polynomials. Singh, Rokne and Dhaliwal [10] have find out the solution of triple series equations involving Laguerre polynomials in a closed form. Kuldeep Narain ([7], [8]), Rajnesh Krishnan Mudaliar and Kuldeep Narain [6] have solved Certain dual and quadruple series equations involving generalized Laguerre polynomials and Jacobi polynomials as kernels. In the present paper, an exact solution has been obtained for the quadruple series equations involving Laguerre polynomials by Noble [9] modified multiplying factor technique.

2010 Mathematics Subject Classifications: 45XX, 33C45, 33D45, 34BXX

Keywords and phrases: Laguerre polynomials, Basic orthogonal polynomials and functions, Boundary value problems.

1 Introduction

Earlier Srivastava ([13], [15]) has solved dual series equations involving Bateman- k functions and Jacobi polynomials. Srivastava [16] has obtained more results like generating functions, bilinear generating functions, recurrence relations, some expansions of functions for the Konhauser-biorhogonal set and general result for the dual series equations involving generalized Laguerre polynomials by putting $k = 1$ in (3.10) and (3.11) in [15,p.645]. Lowndes ([3], [4]), Srivastava [12], Lowndes and Srivastava [5], Srivastava[14], Srivastava and Panda [17] have obtained the solution of dual series equations involving Jacobi and Laguerre polynomials and also solved triple series equations involving Laguerre polynomials. Singh, Rokne and Dhaliwal [10] have find out the solution of triple series equations involving Laguerre polynomials in a closed form. Kuldeep Narain ([7], [8]), Rajnesh Krishnan Mudaliar and Kuldeep Narain [6] have solved Certain dual and quadruple series equations involving generalized Laguerre polynomials and Jacobi polynomials as kernels. In this paper, we have obtained the solution of the following quadruple series equations:

$$(1.1) \sum_{n=0}^{\infty} \frac{A_n}{\Gamma(\alpha+n+p+1)} L_{n+p}^{\alpha}(x) = \phi_1(x), 0 \leq x < a,$$

$$(1.2) \sum_{n=0}^{\infty} \frac{A_n}{\Gamma(\alpha+\beta+n+p)} L_{n+p}^{\alpha}(x) = \phi_2(x), a < x < b,$$

$$(1.3) \sum_{n=0}^{\infty} \frac{A_n}{\Gamma(\alpha+n+p+1)} L_{n+p}^{\alpha}(x) = \phi_3(x), b < x < c,$$

$$(1.4) \sum_{n=0}^{\infty} \frac{A_n}{\Gamma(\alpha+\beta+n+p)} L_{n+p}^{\alpha}(x) = \phi_4(x), c < x < \infty,$$

where $0 < \beta + m$, $0 < \alpha + \beta < \alpha + 1$, p and m are non-negative integer.

$$(1.5) L_{n+p}^{\alpha}(x) = \binom{\alpha+n+p}{n+p}, F_1[-n-p; \alpha+1; x]$$

is the Laguerre Polynomial, $\phi_1(x)$, $\phi_2(x)$, ϕ_3 and $\phi_4(x)$ are prescribed functions.

The solution presented in this paper is obtained by employing a multiplying factor technique similar to that used by Noble [9] or Lowndes ([3],[4]).

2 Preliminaries

The results, which will be required in the course of analysis, are given below for ready reference. From Erdélyi [2], it can be deduced that

$$(2.1) \int_0^y x^\alpha (y-x)^{\beta+m-1} L_{n+p}^\alpha(x) dx = \frac{\Gamma(\beta+m)\Gamma(\alpha+n+p+1)}{\Gamma(\alpha+\beta+m+n+p+2)} y^{\alpha+\beta+m} L_{n+p}^{\alpha+\beta+m}(y)$$

where $0 < y < d$, $-1 < k$, $0 < \beta + m$, and

$$(2.2) \int_y^\infty e^{-x} (x-y)^{-\beta} L_{n+p}^\alpha(x) dx = \Gamma(1-\beta) e^{-y} L_{n+p}^{\alpha+\beta-1}(y)$$

where $d < y < \infty$, $\alpha + 1 > \alpha + \beta > 0$.

From Erdélyi [2], we derive the following orthogonality relation for the Laguerre polynomial:

$$(2.3) \int_y^\infty e^{-x} x^\alpha L_m^\alpha(x) L_n^\alpha(x) dx = \frac{\Gamma(\alpha+n+1)}{n!} \delta_{m,n}$$

where $\alpha > -1$ and $\delta_{m,n}$ is the Kronecker delta.

The differentiation formula:

$$(2.4) \frac{d^{m+1}}{dx^{m+1}} \{x^{\alpha+m+1} L_n^{\alpha+m+1}(x)\} = \frac{\Gamma(\alpha+m+n+2)}{\Gamma(\alpha+n+1)} x^\alpha L_n^\alpha(x)$$

follows from Erdélyi [1].

The analysis here is formal and no attempt has been made to justify the various limiting process.

Making an appeal to the results due to Lowndes ([3], p.123, p.126 eqns (5), (20)), he easily derived ([4], p.168, eqn. (10))

$$(2.5) S(r, x) = (r, x)^\alpha \sum_{n=0}^\infty \frac{\Gamma(n+1)\Gamma(\alpha+\beta+n)}{\{\Gamma(\alpha+1+n)\}^2} L_n(\alpha, x) L_n(\alpha, r),$$

$$(2.6) = \frac{1}{\{\Gamma(1-\beta)\}^2} \int_0^t n(y) (r-y)^{-\beta} (x-y)^{-\beta} dy,$$

$$(2.7) = -\frac{1}{\{\Gamma(1-\beta)\}^2} S_t(r, x),$$

where $\beta < 1$, $\alpha + \beta > 0$, $n(y) = e^y y^{\alpha+\beta-1}$ and $t = \min(r, x)$ of $f(x)$ and $f'(x)$ are continuous in $a \leq x \leq b$ and if $0 < \sigma < 1$, then the solutions of the Abel integral equations

$$(2.8) f(x) = \int_0^x \frac{F(y)}{(x-y)^\sigma} dy,$$

$$(2.9) f(x) = \int_x^b \frac{F(y)}{(y-x)^\sigma} dy,$$

are given by

$$(2.10) F(y) = \frac{\sin \sigma \pi}{\pi} \frac{d}{dy} \int_a^y \frac{f(x)}{(x-y)^{1-\sigma}} dx$$

and

$$(2.11) F(y) = \frac{\sin \sigma \pi}{\pi} \frac{d}{dy} \int_y^b \frac{f(x)}{(x-y)^{1-\sigma}} dx$$

respectively.

3 Solution of the problem

Multiply equation (1.1) by $x^\alpha (y-x)^{\beta+m-1}$, integrate with respect to x over $(0, y)$ and then use (2.1) to obtain

$$(3.1) \sum_{n=0}^\infty \frac{A_n}{\Gamma(\alpha+\beta+m+n+p+2)} y^{\alpha+\beta+m} L_{n+p}^{\alpha+\beta+m} = \frac{1}{\Gamma(\beta+m)} \int_0^y x^\alpha (y-x)^{\beta+m-1} \phi_1(x) dx,$$

where $0 < y < a$, $-1 < \alpha$, $0 < \beta + m$ and m is a non-negative integer.

Differentiate (3.1) $(m+1)$ times with respect to y and use (2.4) to find

$$(3.2) \sum_{n=0}^\infty \frac{A_n}{\Gamma(\alpha+\beta+n+p)} L_{n+p}^{\alpha+\beta-1}(y) = \frac{y^{1-\alpha-\beta}}{\Gamma(\beta+m)} \frac{d^{m+1}}{dy^{m+1}} \int_0^y x^\alpha (y-x)^{\beta+m-1} \phi_1(x) dx,$$

where $0 < y < a$, $-1 < \alpha$, $0 < \beta + m$ and m is a non-negative integer. Again multiply (1.2) by $e^{-x} (x-y)^{-\beta}$, integrate with respect to x over (y, ∞) , then use (2.2) to find

$$(3.3) \sum_{n=0}^\infty \frac{A_n}{\Gamma(\alpha+\beta+n+p)} L_{n+p}^{\alpha+\beta-1}(y) = \frac{e^{-y}}{\Gamma(1-\beta)} \int_y^b (x-y)^{-\beta} e^{-x} \phi_2(x) dx,$$

where $a < y < b$, $\beta < 1$ and $0 < \alpha + \beta$.

Now, multiply equation (1.3) by $x^\alpha (y-x)^{\beta+m-1}$, integrate with respect to x over (a, y) , then use (2.1) to get

$$(3.4) \sum_{n=0}^\infty \frac{A_n}{\Gamma(\alpha+\beta+m+n+p+2)} y^{\alpha+\beta+m} L_{n+p}^{\alpha+\beta+m}(y) = \frac{1}{\Gamma(\beta+m)} \int_b^y x^\alpha (y-x)^{\beta+m-1} \phi_3(x) dx,$$

where $b < y < c$, $-1 < \alpha$, $0 < \beta + m$ and m is a non-negative integer.

Differentiate (3.4), $(m + 1)$ times with respect to y and use (2.4) to find

$$(3.5) \quad \sum_{n=0}^{\infty} \frac{A_n}{\Gamma(\alpha+\beta+n+p)} L_{n+p}^{\alpha+\beta-1}(y) = \frac{y^{1-\alpha-\beta}}{\Gamma(\beta+\alpha+m)} \frac{d^{m+1}}{dy^{m+1}} \int_b^y x^\alpha (y-x)^{\beta+m-1} \phi_3(x) dx,$$

where $b < y < c$, $-1 < \alpha$, $0 < \beta + m$ and m is a non-negative integer.

Now multiply (1.4) by $e^{-x}(x-y)^{-\beta}$, integrate with respect to x over (y, ∞) , then use (2.2) to get

$$(3.6) \quad \sum_{n=0}^{\infty} \frac{A_n}{\Gamma(\alpha+\beta+n+p)} L_{n+p}^{\alpha+\beta-1}(y) = \frac{e^{-y}}{\Gamma(1-\beta)} \int_y^{\infty} (x-y)^{-\beta} e^{-x} \phi_4(x) dx,$$

where $c < y < \infty$, $\beta < 1$ and $0 < \alpha + \beta$.

The left hand sides of equations (3.2), (3.3), (3.5), (3.6) are now identical and hence on using orthogonality relation (2.3), we obtain the solution of equations (1.1), (1.2), (1.3) and (1.4) in the form:

$$(3.7) \quad A_n = \frac{(n+p)!}{\Gamma(\beta+m)} \left[\int_0^a e^{-y} L_{n+p}^{\alpha+\beta-1}(y) F_1(y) dy + \int_b^c e^{-y} L_{n+p}^{\alpha+\beta-1}(y) F_3(y) dy \right] \\ + \frac{(n+p)!}{\Gamma(1-\beta)} \left[\int_a^b y^{\alpha+\beta-1} L_{n+p}^{\alpha+\beta-1}(y) F_2(y) dy + \int_c^{\infty} e^{-y} L_{n+p}^{\alpha+\beta-1}(y) F_4(y) dy \right],$$

where

$$(3.8) \quad F_1(y) = \frac{d^{m+1}}{dy^{m+1}} \int_0^y x^\alpha (y-x)^{\beta+m-1} \phi_1(x) dx,$$

$$(3.9) \quad F_2(y) = \int_y^b (x-y)^{-\beta} e^{-x} \phi_2(x) dx,$$

$$(3.10) \quad F_3(y) = \frac{d^{m+1}}{dy^{m+1}} \int_b^y x^\alpha (y-x)^{\beta+m-1} \phi_3(x) dx,$$

$$(3.11) \quad F_4(y) = \int_y^{\infty} (x-y)^{-\beta} e^{-x} \phi_4(x) dx.$$

The solution of Lowndes equations

$$(3.12) \quad \sum_{n=0}^{\infty} C_n \Gamma(\alpha + \beta + n) L_n^\alpha(x) = \phi_1(x), \quad 0 \leq x < a,$$

$$(3.13) \quad \sum_{n=0}^{\infty} C_n \Gamma(\alpha + 1 + n) L_n^\alpha(x) = \phi_2(x), \quad a < x < b,$$

$$(3.14) \quad \sum_{n=0}^{\infty} C_n \Gamma(\alpha + \beta + n) L_n^\alpha(x) = \phi_3(x), \quad b < x < c,$$

$$(3.15) \quad \sum_{n=0}^{\infty} C_n \Gamma(\alpha + 1 + n) L_n^\alpha(x) = \phi_4(x), \quad c < x < \infty,$$

can be obtained by putting $A_n = C_n \Gamma(\alpha + n + 1) \Gamma(\alpha + \beta + n)$ and $p = 0$ in the solution (3.7).

4 Conclusion

The Laguerre polynomials have been applied by many authors like Lowndes ([3],[4]), Srivastava [12], Srivastava and Panda [17], Lowndes and Srivastava [5], Singh, Rokne and Dhaliwal [10], Kuldeep Narain ([7], [8]), Mudaliar and Kuldeep Narain [6] to solve dual, triple and quadruple series equations. The solution presented in this paper is obtained by employing a multiplying factor technique similar to that used by Noble [9] or Lowndes ([3], [4]). Thus we have obtained an exact solution for the quadruple series equations involving Laguerre polynomials by modified multiplying factor technique Noble [9].

Acknowledgement. The authors are very much thankful to the Editors and Reviewers for their valuable suggestions to bring this paper in its present form.

Dedication This paper is dedicated to Prof. R. C. Singh Chandel on his 75th Birth Anniversary Celebrations for his noteworthy contribution to Mathematical Sciences, *Jñānābha* and VPI continuously since 1971.

References

- [1] A. Erdélyi, *Higher Transcendental Functions*, Vol., **2**, Mc Graw Hill, 1953.
- [2] A. Erdélyi, *Tables of Integrals Transforms*, Vol. **2**, Mc Graw Hill, 1954.
- [3] J. S. Lowndes, Some dual series equations involving Laguerre polynomials, *Pacific J. Math*, **25(1)** (1968), 123-127.
- [4] J. S. Lowndes, Triple series equations involving Laguerre polynomials, *Pacific J. Math*, **29(1)** (1969), 167-173.
- [5] J. S. Lowndes and H. M. Srivastava, Some Triple Series and Triple Integral Equations, *Journal of Mathematical Analysis and Applications*, **150** (1990), 181-187

- [6] Rajnesh Krishnan Mudaliar and Kuldeep Narain, Certain Quadruple Integral equations, *Global journal of pure and Applied Mathematics*, **12(3)** (2016), 2867- 2875.
- [7] Kuldeep Narain, Certain Quadruple series equations, *Scientific Research Journal (SCRJ)*, **1(2)** (2013), 26 -30.
- [8] Kuldeep Narain, Certain Quadruple Series equations with Jacobi polynomials as Kernels, *International Journal of Mathematics Research*, **11(1)** (2019), 55-58.
- [9] B. Noble, Some dual series equations involving Jacobi Polynomials, *Proc. Camb. Phil. Soc.*, **59** (1963), 363-371.
- [10] B. M. Singh, J. Rokne and R. S. Dhaliwal, On closed form solution of triple series equations involving Laguerre polynomials, *Ukrainian Mathematical Journal*, **62(2)** (2010), 259-267.
- [11] J.N. Sneddon and R.P. Srivastava, Dual series relations involving Fourier-Bessel Series, *Proc. Roy. Soc. Edin, A* **66** (1914), 150-160.
- [12] H. M. Srivastava, A Note on Certain Dual Series Equations Involving Laguerre Polynomials, *Pacific J. Math.*, **30(2)** (1969), 525-527.
- [13] H. M. Srivastava, A pair of dual series equations involving generalized Bateman k -functions, *Nederl. Akad. Wetensch. Proc. Ser. A 75 = Indag. Math.*, **34** (1972), 53-61.
- [14] H. M. Srivastava, Certain dual series equations involving Jacobi polynomials. I and II, *Atti Accad. Naz. Lincei Rend. Cl. Sci. Fis. Mat. Natur. (Ser. 8)* **67** (1979), 395-401; *ibid.*, **68** (1980), 34-41.
- [15] H. M. Srivastava, Remarks on certain dual series equations involving the Konhauser biorthogonal polynomials, *J. Math. Phys.*, **23** (1982), 357-357.
- [16] H. M. Srivastava, Some families of dual and triple series equations involving the Konhauser biorthogonal polynomials, *Ganita* (Professor R. P. Agarwal Dedication Volume), **43** (1992), 75-84.
- [17] H. M. Srivastava and R. Panda, A certain class of dual equations involving series of Jacobi and Laguerre polynomials, *Nederl. Akad. Wetensch. Proc. Ser. A 81 = Indag. Math.*, **40** (1978), 502-514.